

Class 26 - October 28
Conditional and Indirect Proof in Predicate Logic (§8.4)

I. A problem arising from using CP and IP in Predicate Logic

With unrestricted CP we could construct the following derivation:

1. $(x)Rx \supset (x)Bx$	Premise
2. Rx	ACP
3. $(x)Rx$	2, UG
4. $(x)Bx$	1, 3, MP
5. Bx	4, UI
6. $Rx \supset Bx$	2-5, CP
7. $(x)(Rx \supset Bx)$	6, UG

This would mean that we could prove that everything red is blue (the conclusion) from ‘If everything is red, then everything is blue’ (the premise).

But that premise can be true while the conclusion is false.

So, the derivation should be invalid.

Moral of the story: we must restrict conditional proof.

The problem is in step 3.

We may not generalize on x within the assumption.

The assumption just means that a random thing is R , not that everything is R .

While variables retain their universal character in a proof, when they are used within an assumption (for CP or IP), they lose that universal character.

It is as if we are saying, “Imagine that some (particular) thing has the property ascribed in the assumption.”

If it follows that the thing in the assumption also has other properties, we may generalize after we’ve discharged, as in line 7.

For, we have not made any specific claims about the thing, outside of the assumption.

The Restriction on CP and IP:

Never UG within an assumption on a variable that is free in the first line of the assumption.

II. Examples of CP and IP in Predicate Logic

One of two typical uses of CP:

- | | |
|-----------------------------------|------------------------|
| 1. $(x)[Ax \supset (Bx \vee Dx)]$ | |
| 2. $(x)\sim Bx$ | $/ (x)(Ax \supset Dx)$ |
| 3. Ay | ACP |
| 4. $Ay \supset (By \vee Dy)$ | 1, UI |
| 5. $By \vee Dy$ | 4, 3, MP |
| 6. $\sim By$ | 2, UI |
| 7. Dy | 5, 6, DS |
| 8. $Ay \supset Dy$ | 3-7, CP |
| 9. $(x)(Ax \supset Dx)$ | 8, UG |

Pick a random object that has property A.

Given any object, if it has A, it provably has D.
Since we are no longer within the scope of the assumption, we may UG.

QED

So, to prove statements of the form $(x)(Px \supset Qx)$:

Assume Px .
Derive Qx .
Discharge $(Px \supset Qx)$.
Then use UG.

Another typical use of CP:

- | | |
|---|---|
| 1. $(x)[Px \supset (Qx \cdot Rx)]$ | |
| 2. $(x)(Rx \supset Sx)$ | $/ (\exists x)Px \supset (\exists x)Sx$ |
| 3. $(\exists x)Px$ | ACP |
| 4. Pa | 3, EI |
| 5. $Pa \supset (Qa \cdot Ra)$ | 1, UI |
| 6. $Qa \cdot Ra$ | 5, 4, MP |
| 7. Ra | 6, Com, Simp |
| 8. $Ra \supset Sa$ | 2, UI |
| 9. Sa | 8, 7, MP |
| 10. $(\exists x)Sx$ | 9, EG |
| 11. $(\exists x)Px \supset (\exists x)Sx$ | 3-10, CP |

QED

Indirect Proof works basically in the same way as in propositional logic.
But the same restriction on CP holds for IP.

Typical use of IP:

1. $(x)[(Ax \vee Bx) \supset Ex]$		
2. $(x)[(Ex \vee Dx) \supset \sim Ax]$	$/ (x)\sim Ax$	
3. $\sim (x)\sim Ax$	AIP	Remember, you're looking for a contradiction.
4. $(\exists x)Ax$	3, CQ	
5. Aa	4, EI	
6. $(Ea \vee Da) \supset \sim Aa$	2, UI	
7. $\sim (Ea \vee Da)$	6, 5, DN, MT	
8. $\sim Ea \cdot \sim Da$	7, DM	
9. $\sim Ea$	8, Simp	
10. $(Aa \vee Ba) \supset Ea$	1, UI	
11. $\sim (Aa \vee Ba)$	10, 9, MT	
12. $\sim Aa \cdot \sim Ba$	11, DM	
13. $\sim Aa$	12, Simp	
14. $Aa \cdot \sim Aa$	5, 13, Conj	
15. $(x)\sim Ax$	3-13, IP, DN	

QED

Note that with CP, sometimes you only assume part of a line, then generalize outside the assumption, but with IP, you almost always assume the negation of the whole conclusion.

III. Exercises. Derive the conclusions of the following arguments:

1. $(x)(Fx \supset Gx)$
 2. $(x)(Fx \supset Hx)$ $/ (x)[Fx \supset (Gx \cdot Hx)]$
1. $(x)(Jx \supset \sim Kx)$ $/ \sim (\exists x)(Jx \cdot Kx)$
1. $(x)(Rx \supset Bx)$ $/ (x)Rx \supset (x)Bx$
1. $(x)(Lx \supset Mx)$
 2. $\sim (\exists x)Lx \supset (\exists x)Mx$ $/ \sim (x)\sim Mx$

Solutions may vary.