

Solutions to the ‘Try these’ examples on each worksheet

Translation key for all problems on all five worksheets:

- a: Aristotle; b: Berkeley; c: *The Critique of Pure Reason*; d: Descartes; e: *The Ethics*; f: Frege;
- g: Heidegger; h: Hume; i: Leibniz; k: Kant; l: Locke; n: Nietzsche; p: Plato; q: *The Inquiry Concerning Human Understanding*; r: Arendt; s: Spinoza; t: Socrates
- Bx: x is a book; Cx: x is a coherentist; Ex: x is an empiricist; Ix: x is an idealist; Mx: x is a materialist; Px: x is a philosopher; Rx: x is a rationalist
- Bxy: x is bigger than y; Dxy: x is more difficult to read than y; Lxy: x likes y; Mxy: x is read more widely than y; Oxy: x is more original than y; Pxy: x plays billiards with y; Rxy: x respects y; Sxy: x studies y; Wxy: x wrote y
- Lxyz: x likes y better than z

Only

6. $P_n \bullet M_{nd} \bullet (\forall x)[(Px \bullet M_{xd}) \supset x=n]$
7. $E_k \bullet R_k \bullet (\forall x)[(Ex \bullet Rx) \supset x=k]$
8. $E_l \bullet P_l \bullet (\exists x)(Rx \bullet Px \bullet Rx_l) \bullet E_b \bullet P_b \bullet (\exists x)(Rx \bullet Px \bullet Rx_b) \bullet (\forall x)\{[Ex \bullet Px \bullet (\exists y)(Ry \bullet Py \bullet Ry_x)] \supset (x=l \vee x=b)\}$

Except

6. $P_i \bullet P_b \bullet (\forall x)[(Px \bullet x \neq i \bullet x \neq b) \supset M_x]$
7. $P_r \bullet R_{rg} \bullet (\forall x)[(Px \bullet x \neq r) \supset \sim Rx_g]$
8. $(\exists x)\{Bx \bullet \sim Sn_x \bullet (\forall y)[(Py \bullet y \neq n) \supset Sy_x]\}$

Superlatives

6. $P_s \bullet (\forall x)[(Px \bullet x \neq s) \supset Os_x]$
7. $B_c \bullet W_{kc} \bullet (\forall x)[(Bx \bullet W_{kx} \bullet x \neq c) \supset Mc_x]$
8. $(\exists x)\{Bx \bullet (\exists y)(Ey \bullet Wy_x) \bullet (\forall z)\{[(Bz \bullet (\exists w)(Ew \bullet W_{wz}) \bullet z \neq x) \supset Bx_z]\}\}$

At least

6. $(\exists x)(\exists y)(Px \bullet Py \bullet M_{xf} \bullet My_f \bullet x \neq y)$
7. $(\exists x)(\exists y)(\exists z)(Px \bullet Py \bullet Pz \bullet M_{xf} \bullet My_f \bullet M_{zf} \bullet x \neq y \bullet x \neq z \bullet y \neq z)$
8. $(\exists x)(\exists y)(\exists z)(\exists w)(Ix \bullet Iy \bullet Iz \bullet Iw \bullet S_{xc} \bullet S_{yc} \bullet S_{zc} \bullet S_{wc} \bullet x \neq y \bullet x \neq z \bullet x \neq w \bullet y \neq z \bullet y \neq w \bullet z \neq w)$

At most

6. $(\forall x)(\forall y)[(Px \bullet Ex \bullet Rx \bullet Py \bullet Ey \bullet Ry \bullet Pz \bullet Ez \bullet Rz) \supset x=y]$
7. $(\forall x)(\forall y)(\forall z)[(Px \bullet R_{bx} \bullet Py \bullet R_{by} \bullet Pz \bullet R_{bz}) \supset (x=y \vee x=z \vee y=z)]$
8. $(\exists x)(Ex \bullet L_{xd}) \bullet (\forall x)(\forall y)(\forall z)[(Ex \bullet L_{xd} \bullet Ey \bullet Ly_d \bullet Ez \bullet L_{zd}) \supset (x=y \vee x=z \vee y=z)]$